

Froduct Name:	FIOLEIII MULLICOIOI Stable II	
Code No:	DM660	
Lot No:	018CB11	
Size:	1.2 ml (120 mini-gel lanes)	
Storage:	4 °C	
Stability:	12 months at 4 °C	
Storage Buffer:	Tris-HCl (pH6.8), EDTA, Glycerol, SDS	

# Product Name: DynaMarker Protein MultiColor Stable II

### Description

The <sup>DynaMarker</sup> Protein MultiColor Stable II is a pre-stained protein molecular weight marker. The marker has a remarkable feature that it is possible to store at 4 °C. The feature allow us to start electrophoresis with the marker, because it is always in a liquid state while stored at 4 °C. The <sup>DynaMarker</sup> Protein MultiColor Stable II consists of nine prestained proteins. Each of them are stained red, blue, purple, green or orange, ranging in apparent molecular weight from approximately 8 kDa to 230 kDa. The <sup>DynaMarker</sup> Protein MultiColor Stable II is suitable for visualizing proteins during electrophoresis without staining and for monitoring electrophoretic transfer onto membranes. The protein concentrations are optimized to give uniform band intensities. The marker is supplied in gel loading buffer for direct loading onto SDS-PAGE without heating or adding reducing agents.



Electrophoresis profile of <sup>DynaMarker</sup> Protein MultiColor Stable II (10 µl) on 6% polyacrylamide (5% C) Gel / AllView PAGE Buffer (Code#DS520) as running buffer.

## Protocol

- 1. Take the marker out of refrigerator.
- 2. Load 10  $\mu$ l for mini-gels or more for large size gels.
- 3. Load your samples.
- 4. Start electrophoresis.

Note: There is no need to heat or add reducing agents.



# Contents

	Apparent molecular weight (kDa) *		
	Tris-Glycine-SDS buffer	AllView PAGE Buffer	
Protein	(Laemmli running buffer)	(Code#DS520)	
Myosin	249	250	
β-Galactosidase	137	141	
Phosphorylase-b	96	100	
BSA	73	71	
Ovalbumin	46	45	
Carbonic anhydrase	31	30	
Soybean trypsin inhibitor	26	25	
Lysozyme	18	17	
Aprotinin	6.3	8.4	

**Note:** As covalently bound dye affects protein mobility, each batch of prestained protein marker is calibrated against unstained standards. A prestained protein marker should be used for approximate molecular weight determination. For precise molecular weight determination use an unstained molecular weight marker.

\*: The apparent molecular weight values are lot specific and depends on the electrophoresis running buffer.

Code	Product Name	Description
DS520	AllView PAGE Buffer	AllView PAGE Buffer is a new type of running buffer
		for SDS-PAGE electrophoresis. This buffer is possible
		to separate proteins with wide range of molecular
		weights in the basic Laemmli gel (Tris-HCl) similar to
		using "gradient gel".

### **Related Products**